

Joint Injection Techniques



Brian Coleman, M.D.
OU Family Medicine
23 SEP 2025

Learning Objectives

- To describe the benefits and risks of joint injection
 - To identify situations when joint injections should not be performed
 - To identify the unique anatomy of commonly injected joints which allows for successful joint injection
- To demonstrate the appropriate techniques for performing injection of shoulder and knee



General Principles

- Anatomical placement
- Local anesthetic blocks
- Corticosteroid injection vs visco-supplementation vs regenerative (PRP/stem cells/etc)
- Rest/restricted use
- Adjunctive physical therapy




Inject with caution

- Patellar tendon
- Achilles tendon
- Biceps tendon
- Infected joint



Materials

- **Prep materials**
 - Betadine, alcohol preps, chlorhexidine, chloraprep
- **Anesthetic**
 - Ethyl chloride
 - 1% lidocaine without epinephrine
- **Steroid of choice**
- **Syringe**




Materials

- Needles
- Medium hemostat
- Post-injection prep
 - Band-aids, 4 x 4 gauze, 1" tape, elastic wrap
- Epinephrine readily available




Benefits to injection

- Pain relief, improved function
 - Viscosupplementation--effective treatment for knee OA of the knee—improved pain, function and patient global assessment (Cochrane meta-analysis)
 - Insufficient evidence with corticosteroid injections (Cochrane database)
 - Lidocaine injection into myofascial trigger points appears effective (Cochrane database)




Absolute Contraindications

- Local cellulitis
- Joint prosthesis
- Septic arthritis
- Bacteremia
- Acute fracture
- History of allergy or anaphylaxis to injectable pharmaceuticals or constituents



Relative contraindications


- Minimal relief after two previous corticosteroid injections
- Underlying coagulopathy
- Anticoagulation therapy
- Evidence of surrounding joint osteoporosis
- Anatomically inaccessible joints
- Uncontrolled diabetes mellitus



Side Effects

All patients should be warned about:


- 30%--soreness or pain after injection (2-3 days, ice, Tylenol)
- 10%--inflammatory flare reaction (2-3 days, ice, narcotics)
- 30%--fat or skin atrophy (90% revert to normal in 6-12 months)
- Reevaluate if redness, swelling, and pain beyond 3-4 days (infection risk < 1:10,000)



Side Effects

Remember these complications also:

- **Tendon rupture (if injected into the tendon)**
- **Damage to the cartilage after repeated injection**
- **Can cause a crystalline arthritis**



Synovial Fluid

- Can't be aspirated from a healthy joint
- Often holds key to diagnosis
- Normally, fluid is transparent, doesn't contain large proteins or clot
- Send fluid for analysis, cultures, protein, cell count, and crystals



Synovial Fluid Analysis

Parameter	Normal	Noninflammatory	Inflammatory	Septic
Clarity	Transparent	Transparent	Translucent to opaque	Opaque
Color	Clear	Yellow	Yellow to green	Yellow to green
Viscosity	High	High	Low	Variable
WBC/mm ³	<200	200-2,000	2,000-150,000	15,000-200,000
Polys	<25%	<25%	>50%	>75%
Culture	Negative	Negative	Negative	Usually Positive
Protein (gm/dl)	<2.5	<2.5	>2.5	>2.5

Common Injectable Corticosteroids

Medication	Potency	Onset	Duration
Hydrocortisone (cortisol)	1	Fast	Short
Prednisolone terbutate (Hydeltra)	4	Fast	Intermediate
Methylprednisolone acetate (Depo-Medrol)	4	Slow	Intermediate
Triamcinolone acetonide (Kenalog)	5	Moderate	Intermediate
Triamcinolone hexacetonide (Aristospan)	5	Moderate	Intermediate
Betamethasone (Celestone)	25	Fast	Long
Dexamethasone sodium phosphate (Decadron)	25	Fast	Long

Usual Doses of Methylprednisolone or Equivalent by Site

Dose	Anatomic site
5 to 10 mg	Phalangeal joints
20 to 30 mg	Wrist
20 to 30 mg	Elbow and ankle
40 to 80 mg	Shoulder, hip, or knee



Common Injection Sites

- Shoulders
- Elbow
- Hand and wrist
- Hip
- Knee
- Foot and ankle
- Trigger points



Shoulder


Three locations

- Subacromial/subdeltoid bursa
 - Impingement, rotator cuff tear
- Acromioclavicular joint
 - osteoarthritis
- Glenohumeral joint




Subacromial Bursa

- Palpate the acromion and posterior humeral head
- Use 1.5" 25 gauge needle, instilling 10 cc of steroid/anesthetic combo
- Enter posteriorly aiming toward the coracoid process anteriorly
- Enter laterally, pulling down on the arm to open the joint up, aiming slightly upward under the acromion



**Subacromial
bursa**








Figure 1 consists of two panels. Panel (a) is a clinical photograph showing a patient's shoulder with a needle inserted into the subacromial space. Panel (b) is an ultrasound image of the shoulder joint, showing the needle tip and the subacromial space. The image includes labels for the needle, subacromial space, and biceps tendon. The ultrasound image also displays technical parameters on the right side, such as frequency (7.0 MHz), depth (3.0 cm), and gain (100%).



Acromioclavicular Joint

- Palpate distal edge of clavicle, proceed until you feel a bump (this is the joint)
- Insert 1" 25 gauge needle from posterior surface, angle medially (will feel needle slip into place)
- Use small volume, 2-5 ml of steroid-anesthetic combination






Elbow

Common reasons for injection/aspiration of the elbow:

- **Lateral/medial epicondylitis**
- **Olecranon bursitis**
- **Degenerative arthritis**




Epicondylitis

- **Locate most tender area. Insert 1.5" 25 gauge needle at 90° angle, go down to the periosteum and begin injecting into it (key to a good response), withdraw slightly and complete infiltration**
 - **Be careful to not infiltrate the ulnar or radial nerve**
 - **Massage medication into the region to disperse**
- Rest for 7-10 days**




LATERAL EPICONDYLE



Olecranon Bursitis



- Elbow 90° of flexion, insert the needle from the back of the elbow, parallel to ulna posterior
- Use larger needle (20 or 22 gauge)
- If fluid is clear, then may use steroids; but if slightly cloudy, don't





DeQuervain's Tenosynovitis

- Insert 1" 25 gauge needle over the tendon sheath at 45° angle
- Should balloon/dilate as infiltrates, signifies medication is in sheath




Abductor
pollicis
longus



Trigger Finger


- Locate nodule by palpating tendon at junction of A-1 pulley, distal to the distal palmar crease
- Tendon is very superficial (2-3 mm) deep
- Inject on top/volar surface of flexor tendon
- 50% resolve within days






Carpal Tunnel Syndrome



- Locate palmaris longus tendon at proximal crease of wrist
- Median nerve is directly under tendon
- Insert 1.5" 25 gauge needle just ulnar to tendon using 45° angle, aiming distally
- Begin injecting when 1" deep
If feels pain or tingling, need to reposition needle





← Palmaris longus


← Injection site





Knee


Two Common Methods

- **Knee extended**
 - Palpate superior pole of patella
 - Insert needle laterally/medially just deep to patella into suprapatellar pouch
- **Knee flexed**
 - Insert needle at the level of inferior pole, lateral/medial to the patellar ligament into intercondylar notch
- **Aspiration = 18-20 gauge 1.5"**
- **Injection = 22 gauge 1.5" needle**











Hip

- Intra-articular joint injection → orthopedic referral—use guidance for injection
- Greater trochanteric bursitis
 - Patient lay on their side with affected side up
 - Palpate area of maximal tenderness
 - Insert 3.5” 25 gauge spinal needle down to periosteum, infiltrate some on periosteum and withdraw slightly to complete



Trochanteric bursa injection




© NMEG 2008



Ankle

- Aspiration = 1.5" 18 gauge;
Injection = 1.5" 25 gauge needle
- Insert needle 0.5 cm lateral to medial malleolus and medial to extensor hallucis longus tendon
- Direct the needle toward dome of the talus






Ankle

Subtalar Arthritis

- Inject into the sinus tarsi on the lateral aspect of the ankle, just below the lateral malleolus
- Inverting the foot will help to locate the space
- Use a 1½" 25 gauge, direct needle posterior and medially, as you walk off the talus of calcaneus

Can use 5-10 ml anesthetic-steroid mixture





Foot

Plantar Fasciitis and Heel Spurs

- Locate point of maximal tenderness on plantar heel, go deep to the bone
- Avoid fat pad as can cause fat atrophy





Morton's Neuroma




Neuroma

Diana Vashilova



Trigger points



Coding issues

- 20600 Arthrocentesis, aspiration and/or injection. small joint or bursa. (eg, finger, toes)
- 20611 – injection large joint using US guidance.
- 20605 intermediate joint or bursa (eg, temporomandibular, acromioclavicular, wrist, elbow or ankle)
- 20610 major joint or bursa (eg, shoulder, hip, knee)
- 20526 Injection, therapeutic (eg. local anesthetic, corticosteroid), carpal tunnel.



Coding Issues (cont)

- 20550 Injection(s), single tendon sheath, or ligament, aponeurosis (eg, plantar "fascia")
- 20551 single tendon origin/insertion
- 20552 Injection(s), single or multiple trigger point(s)
- 20553 single or multiple trigger point(s), three or more muscle(s)



Conclusions

- Understand risks and benefits of aspiration and injections
- Anatomic placement important
- Easy procedure to add to your armamentarium



THE END----- questions?

